

Measure for Measure: Prescribing Guidelines, Rules & Regulations

Stephen J. Ziegler, PhD, JD

Disclosures

- Stephen J. Ziegler, PhD, JD
 - -Nothing to disclose



Learning Objectives

- Discuss the diffusion and variation in prescribing guidelines and rules across the United States regarding chronic, noncancer pain
- Describe the history of Washington state's prescribing rules and the CDC's prescribing guidelines
- Discuss the potential impact that prescribing guidelines could have on medical practice, the treatment of pain, and reduction of overdose




Increases in Overdose

Drug overdose death rates in the United States have never been higher

- Drug overdose death rates have risen steadily in the United States since 1970. (See Figure 1)
- In 2007, 27,658 unintentional drug overdose deaths occurred in the United States.
- Drug overdose deaths were second only to motor vehicle crash deaths among leading causes of unintentional injury death in 2007 in the United States.

Rates have increased roughly five-fold since 1990.


 Age-adjusted rates of drug overdose death for whites have exceeded those among African Americans since 2003.





Some Variables Associated With Overdose Risk

OD Variables Not Equally Distributed



- Illicit opioids
- □ Poly-pharm (Benzo)
- Poly-substance
- Unauthorized >
- Drug interactions
- □ Genotype
- **■** Methadone
- Depression

Painweek.

Hydraulic Pressure to do Something

- Great cases, like hard cases, make bad law. For great cases are called great not by reason of their real importance in shaping the law of the future, <u>but because of some accident of immediate</u> <u>overwhelming interest which appeals to the feelings and distorts the</u> judgment.
- These immediate interests exercise a kind of hydraulic pressure which makes what was previously clear seem doubtful, and by which even well settled principles of law will bend.
- [emphasis added] Justice Holmes (dissenting in Northern Securities Co. v. United States, 193 U.S. 197 (1904))
- Korematsu v. United States, 323 U.S. 214 (1944)



Do Something =

- Hysteria
- Simplistic explanations, bumper sticker solutions, sacred cows
- Result: Rx guidelines, rules, and regulations
- Do the terms get mixed up?
 - -Voluntary guideline vs mandatory rule
 - -Language matters: shall vs should
- Original purpose of <u>guidelines</u>: support clinical decision making
- But expanding ... (regulators, insurance, *standards*?)



7

Definitions: Guidelines vs Rules

- Washington State DOH
- Guidelines
- A set of <u>recommended</u> practices designed by the Medical Commission to <u>assist</u> practitioners about appropriate health care for specific circumstances
- A guideline does <u>not</u> have the force of law, but <u>may</u> be considered by the Medical Commission to be the <u>standard</u> <u>of care in our state</u>. [Emphasis not in original]



В

Guideline or Rule?



Interagency Guideline on Prescribing Opioids for Pain

Developed by the Washington State Agency Medical Directors' Group
(AMDG) in collaboration with an Expert Advisory Panel, Actively Practicing



Federation of State Medical Boards

- Guidelines for the Chronic Use of Opioid Analgesics
- "specific limitation and restriction that these Guidelines do not operate to create any specific standard of care, which standard must depend upon fact-specific totality of circumstances surrounding specific quality of care events."
- The Guidelines recognize that there is not just one appropriate strategy to accomplish the goals of these Guidelines. Effective means of achieving the goals of these Guidelines vary widely depending on the type and causes of the patient's pain, the preferences of the clinician and the patient, the resources available at the time of care, and other concurrent issues beyond the scope of these Guidelines.
- These Guidelines that follow do not encourage the prescribing of opioids over other pharmacological and nonpharmacological means of treatment but rather the Guidelines recognize the responsibility of clinicians to view pain management as essential to quality of medical practice and to the quality of life for patients who suffer from pain."



CDC = Guideline?

- "The recommendations in the guideline are voluntary, rather than prescriptive standards.
- They are based on emerging evidence, including observational studies or randomized clinical trials with notable limitations.
- Clinicians should consider the circumstances and unique needs of each patient when providing care."



But "Rules and Regulations" are different from Guidelines

- Carry force and effect of law
- Created by
- Admin/exec agencies who provide the clothing
- Open comment period
- Testimony (written, oral)
- Opportunity to speak!





Regulations are assigned a number

- Published and codified = assigned administrative law chapter and section number (WAC)
- Many states have such regulations (that can be referred to as rules)

PAIN MANAGEMENT

246-919-850 Pain management—intent.
246-919-851 Exclusions.
246-919-852 Definitions.
246-919-853 Patient evaluation.
246-919-854 Treatment plan.
246-919-855 Informed consent.
246-919-857 Periodic review.
246-919-857 Periodic review.
246-919-858 Episodic care.
246-919-860 Consultation—Recommendations and requirements.
246-919-860 Consultation—Exemptions for exigent and special circumstances.
246-919-863 Pain management specialist.



Common in Guidelines and Rules

- Patient eval, tx agreements, follow-up, check PMDP . . .
- But dosage thresholds (leading to trigger or ceiling) expanding across US
 - -Tied to MED—the myth of MED and its dangers (Fudin et al, 2016)



Indiana (your state med Assn more helpful)

- 844 IAC 5-6-3 Triggers for imposition of requirements; exemptions
- Sec. 3. (a) This section and sections 4 through 10 of this rule establish requirements concerning the use of opioids for chronic pain management for patients.
- (b) . . . this section . . . shall not apply to the use of opioids for chronic pain management for the following:
- (1) Patients with a terminal condition.
- (2) Residents of a health facility licensed under IC 16-28.
- (3) Patients enrolled in a hospice program licensed under IC 16-25.
- (4) Patients enrolled in an inpatient or outpatient palliative care program of a hospital licensed under IC 16-21 or a hospice licensed under IC 16-25.



Indiana Triggers

- (c) The requirements in the sections identified in subsection (a) only apply if a patient has been prescribed [for more than 3 consecutive months]:
- (1) more than sixty (60) opioid-containing pills a month;
- (2) a morphine equivalent dose of more than fifteen (15) milligrams per day;
- (3) a transdermal opioid patch;
- (4) at any time it is classified as a controlled substance under Indiana law, tramadol, but only if the patient's tramadol dose reaches a morphine equivalent dose of more than sixty (60) milligrams per day
- or
- (5) an extended release opioid medication that is not in an abuse deterrent form for which an FDA-approved abuse deterrent formulation, if such a formulation exists for the opioid product the physician is prescribing to the patient. Nothing in this subdivision shall be construed to require a physician to prescribe an opioid in an abuse deterrent formulation.



Origin of Dosage Thresholds



Painweek.

Washington State's History

- Educational pilot
- ■MED 120
- Guideline (not a mandatory rule)
- ■CME?
- Rule



Where did 120 come from?

- "there does not seem to be an evidence-based threshold for what constitutes a dangerously high dose. Although some clinical guidelines suggest an MED of 200 milligrams per day as a watchful dose, studies in our sample showed overdose and mortality increases at doses ranging from 40 to 200 milligrams per day MED." (King, et al, 2014) [systematic review of studies 1990-2013]
- "Ecologic studies suggest a near-linear association between the total amount of opioids dispensed and overdose morbidity and mortality . . . but there is new evidence that the shape of the curve is not linear. (Dasgupta et al, 2016)
- Higher doses of opioid analgesics were associated with increased overdose risk, however, there were smaller incremental increases in risk above 200 mg average daily MME. <u>Much of the risk at higher doses appears to be associated with coprescribed benzodiazepines</u>" (Dasgupta et al, 2016)



Dose alone not problem

- "Although risk for overdose was highest in those receiving higher doses, most overdoses occurred in patients receiving low- to moderate-dose regimens" (Dunn et. al, 2010 at 90). [emphasis added]
- "However, there does not seem to be an evidence based threshold for what constitutes a dangerously high dose. Although some clinical guidelines suggest an MED of 200 milligrams per day as a "watchful dose," studies in our sample showed overdose and mortality increases at doses ranging from 40 to 200 milligrams per day MED." (King et. al, 2014 at e35).



So guidelines may not be the panacea?

- "Many poisonings and adverse effects occurred in patients without high dose or long-term opioid therapy,
- suggesting that opioid dosing and duration guidelines may not be sufficient to reduce morbidity related to prescription opioid use.


Fulton-Kehoe et.al, 2013, at 1452)." [emphasis added]




Diffusion, Implementation, & Complication across the US









Laboratories of the State?

- Different MED by state? Medicaid vs non?
- Calculation & conversion: something lost in translation.
- Washington state: CAUTION: This calculator should NOT be used to determine doses when converting a patient from one opioid to another. This is especially important for fentanyl and methadone conversions. Equianalgesic dose ratios are only approximations and do not account for genetic factors, incomplete cross-tolerance, and pharmacokinetics. [Recall earlier diagram about variables associated with OD?]



MED, CDC, Rx w/ CONFIDENCE! Opioid Overdose CDC > Opioid Overdose > Information for Providers > Guideline Resources Opioid Overdose Opioid Basics Guideline Resources: CDC Opioid Guideline Mobile App Overdose Prevention Information for Patients CDC's new Opioid Guideline App is designed to help providers apply the recommendations of CDC's CDC Opioid Guideline App: Guideline for Prescribing Opioids for Chronic Pain into clinical practice by putting the entire guideline. Prescribe with Confidence tools, and resources in the palm of their hand. Managing chronic pain is complex, but accessing Guideline Overview prescribing guidance has never been easier. The application includes a Morphine Milligram Equivalent (MME) calculator*, summaries of key Clinical Tools $help\ providers\ practice\ effective\ communications\ skills\ and\ prescribe\ with\ confidence.$ Videos The new CDC Opioid Guideline App is now available for free download on Google Play & (Android Mobile App devices) and in the Apple Store of (iOS devices). Training for Providers Painweek.

BEWARE OF THE SMALL PRINT

- What a difference an * makes
- The calculator does not account for incomplete crosstolerance between opioids and should not be used to guide opioid rotation or conversion between different opioids.
- This is especially important for fentanyl and methadone conversions.
- Equianalgesic dose ratios are approximations and do not account for interactions between opioids and other drugs, patient weight, hepatic or renal insufficiency, genetic factors, and other factors affecting pharmacokinetics.
- https://www.cdc.gov/drugoverdose/prescribing/app.html



Problems with primary fixation on daily MED

- Dosage is important
- But multiple contributors to OD
- Association not = causation
- Conversion tables = under and overdosing
 - -"An increasing body of literature suggests that widely used opioid rotation practices, including the use of dose conversion ratios found in equianalgesic tables, may be an important contributor to the increasing incidence of opioid-related fatalities." Webster & Fine (2012).
- Race to bottom in thresholds by states?
- Methadone & Seattle Times (Berens & Armstrong, 2011)



Impact on Patients and Prescribers: Opting Out

Wednesday, October 19, 2016

TO: Washington State Medical Association, Washington State Hospital Association FROM: Melanie de Leon, JD, Executive Director

SUBJECT: Technical Assistance Regarding Noncancer Pain Rules

At the request of several organizations in Washington, we are restating points around the chronic noncancer pain rules (pain rules) effective in 2012. It is our sincere hope that this clarification will promote understanding among the practitioner community and encourage more acceptances of patients dealing with chronic noncancer pain. It is our further hope that this clarification will encourage more practitioners, outside the realm of pain management specialists (e.g., primary care providers), to accept these patients into their practice in good faith and deliver the competent care, including pain care, that we know they are qualified to provide.

1. The pain rules are not intended to force practitioners who are not pain management specialists away from treating chronic noncancer pain patients. In fact the intent is the complete opposite. The goal of the Commission in implementing the pain rules is to assist practitioners by articulating in rule what was previously required by the unwritten standard of care.



Opioid Refugees?

- Many patients have functioned well for years
- Involuntary tapers & "detox"
- Maine: 100. Without consent. Last minute save undermines the rule.
- LTD alternatives effective & reimbursable
- Self-medication
- Only 12% med boards have action plan to respond to loss of CS prescriber (Sera et al, 2017)



The Future?

- Thresholds because they are so easy? Race to bottom
- Pendulum swing & Other Side of Opioids series
- Alternatives: New Mexico CME, Project Lazarus
- CARA (2016)
- Become part of solution
 - Advocate for balanced policies (guidelines, laws, and REGULATIONS—testify)
 - -> Reimbursement>
 - Common ground





Contact Information

Stephen J. Ziegler, PhD, JD Professor Emeritus, Purdue University, Fort Wayne Sziegler@Purdue.edu



References

- Albert S, Brason FW 2nd, Sanford CK, et al. Project Lazarus: Community-based overdose prevention in rural North Carolina. Pain Med 2011; 12 (suppl 2):S77–85.
- Berens MJ, Armstrong K: State pushes prescription painkiller methadone, saving millions but costing lives. Available at: http://bit.ly/2tsPmZs (last accessed June 30, 2017)
- CARA (Comprehensive Addiction & Recovery Act) (2016) https://www.congress.gov/bill/114th-congress/senate-bill/524/text
- Centers for Disease Control and Prevention. Prescription Pain Killer Overdoses, Use and Abuse of Methadone as a painkiller. Vital Signs, July 2012. http://www.cdc.gov/vitalSigns/MethadoneOverdoses/index.html
- Dasgupta N, Funk MJ, Proescholdbell S, et al. Cohort Study of the Impact of High-Dose Opioid Analgesics on Overdose Mortality. Pain Med 2016; 17: 85–98.
- Dunn KM, Saunders KW, Rutter CM, et al. Opioid prescriptions for chronic pain and overdose: A cohort study. Ann Intern Med 2010;152:85–92.



References

- Fishman S. The Politics of Pain and its Impact on Pain Medicine. Pain Med, 2005; 6(3), 199-200.
- Fudin J, Marcoux MD, Fudin JA. Mathematical Model For Methadone Conversion Examined. Practical Pain Management. Sept. 2012. 46-51.
- Fudin J, Cleary JP, Schatman, ME. The MEDD myth: the impact of pseudoscience on pain research and prescribing-guideline development. J Pain Res. 2016; 9: 153–156.
- Greenfield, S. Clinical Practice Guidelines, Expanded Use and Misuse. JAMA. 2017; 317 (6), 594-95.
- Hawryluk M. Opioid Crisis: Pain patients pushed to the brink. Bend Bulletin, June 2, 2017. http://bit.ly/2ruqxur (last accessed June 5, 2017).
- Katzman JG, Comerci GD, Landen M, et al.. Rules and Values: A Coordinated Regulatory and Educational Approach to the Public Health Crises of Chronic Pain and Addiction. Am J Pub Health 2014; 104(8):1356-62.



References

- King NB, Fraser V, Boikos C, Richardson R Harper S. Determinants of increased opioid-related mortality in the United States and Canada, 1990–2013: A systematic review. Am J Pub Health 2014;104:e32–42.
- Merton RK. The Unanticipated Consequences of Purposive Social Action. American Soc Rev. 1936; 1 (6), 894-904, 898, 901.
- Sera L, Brown M, McPherson ML, et al. JOM, 2017; 13 (2), 105-110.
- Shaw K, Fudin J. Evaluation and Comparison of Online Equianalgesic Opioid Dose Conversion Calculators. PPM 2013 (August);13(7):61-66.
- Twillman B. What's Really Driving Opioid-Related Death Rates? Pain Treatment Topics, Jan. 1, 2012, available at: http://updates.pain-topics.org/2012/01/whats-really-driving-opioid-related.html
- Webster LR, Cochella S, Dasgupta N, et al. An Analysis of the Root Causes for Opioid-Related Overdose Deaths in the United States. Pain Med 2011; 12, S26–S35.
- Webster LR, Dasgupta N. Obtaining Adequate Data to Determine Causes of Opioid-Related Overdose Deaths. Pain Med 2011; 12, S86–S92.



References

- Webster LR, Fine PG. Review and Critique of Opioid Rotation Practices and Associated Risks of Toxicity. Pain Med 2012; 13: 562–570.
- Ziegler SJ. The proliferation of dosage thresholds in opioid prescribing policies and their potential to increase pain and opioid-related mortality. Pain Med, 2015; 16 (10), 1851-1856.
- Ziegler SJ. Patient Abandonment in the Name of Opioid Safety. Pain Med, 2013; 14 (3), 323-324.
- Ziegler SJ. Governmental Intervention in Prescribing: Reducing Harm and Medicolegal latrogenesis, PWJ, 2013; 1(3), 36-40.
- Wildland fire photograph. http://bit.ly/2tedg9Y (last accessed July 4, 2017).
- Opioid Dosage and Morphine Equivalency: Implications for Meeting the Standard of Care When Comparing CDC Guidelines to State Guidelines (An Educational Resource for Clinicians), Academy of Integrative Pain Management, Sonora, CA. http://sppan.aapainmanage.org/ (last accessed July 4, 2017).



References

- Links to the Centers for Disease Control and Prevention and state prescribing guidelines & rules (Last accessed June 30, 2017):
- CDC Prescribing Guideline: Dowell D, Haegerich TM, Chou R. CDC Guideline for Prescribing Opioids for Chronic Pain — United States, 2016. MMWR Recomm Rep 2016;65(No. RR-1):1– 49. DOI: http://dx.doi.org/10.15585/mmwr.rr6501e1.
- Guidelines for the Chronic Use of Opioid Analgesics. Federation of State Medical Boards. April 2017. http://bit.ly/2GdgmA5
- Alabama. Alabama State Board of Medical Examiners Rules & Regulations, http://bit.ly/2t9kXg2
- California. Guidelines for prescribing controlled substances for pain, Medical Board of California. http://bit.ly/2ussJBd
- Colorado. Open Letter to the General Public on the Quad-Regulator Joint Policy for Prescribing and Dispensing Opioids. http://bit.ly/2g1IM27
- Indiana. Indiana Pain Management Prescribing Final Rule. http://bit.ly/2tuSJ1o
- Maine. An Act to Prevent Opiate Abuse by Strengthening the Controlled Substances Prescription Monitoring Program. http://bit.ly/2sZULWJ



References

- Massachusetts: Commonwealth of Massachusetts Department of Industrial Accidents, Treatment Guideline 27 Chronic Pain. http://bit.ly/2sZKhGP
- Minnesota. Minnesota Department of Human Services. http://bit.ly/1CuiL2Y
- New Hampshire. Office of Professional Licensure and Certification. http://bit.lv/2ttCtOG
- Ohio. State Medical Board of Ohio, Prescriber Resources. http://bit.ly/2sZCNDK
- Rhode Island. Rules and Regulations for Pain Management. http://bit.ly/2n441aV
- South Carolina. Joint Revised Pain Management Guidelines Approved by the South Carolina Boards of Medical Examiners. http://bit.ly/2sZOFWj
- Vermont. Vermont Department of Health. http://bit.ly/2sZX9No
- Virginia. Regulations Governing Prescribing of Opioids and Buprenorphine. http://bit.ly/2sv8iCr
- Washington state. Medical Quality Assurance Commission. http://bit.ly/2ty5Ek9
- The Other Side of Opioids. http://www.lasvegasnow.com/news/ourpain-the-other-side-of-opioids/852432872

